

Local Realistic Model for the Dynamics of Bulk-Ensemble NMR Information Processing

N. C. Menicucci and Carlton M. Caves

Department of Physics and Astronomy, University of New Mexico, Albuquerque, NM 87131-1156

(November 19, 2001)

Abstract

We construct a local realistic hidden-variable model that describes the states and dynamics of bulk-ensemble NMR information processing up to about 12 nuclear spins. The existence of such a model rules out violation of any Bell inequality, temporal or otherwise, in present high-temperature, liquid-state NMR experiments. The model does not provide an efficient description in that the number of hidden variables grows exponentially with the number of nuclear spins.

High-temperature, liquid-state nuclear magnetic resonance (NMR) provides a testing ground for the new ideas for information processing that are being developed in quantum information science [1]. The fundamental information-processing elements used in NMR are two-level nuclear spins, called *qubits*, which are bound together in a single molecule. A liquid NMR sample contains a macroscopic number of molecules, each of which functions as an independent information-processing unit. The molecules are initially in thermal equilibrium at high enough temperature that the nuclear spins are only weakly polarized along the direction of a strong magnetic field. NMR techniques cannot control the quantum states of individual molecules, and the measurements performed in NMR detect the average magnetization of the entire sample. For these reasons the use of high-temperature, liquid-state NMR to emulate quantum computation is called *bulk-ensemble quantum computation*.

The original proposals [2,3] for quantum information processing using NMR were greeted with enthusiasm tempered by skepticism. The enthusiasm led to a remarkable series of experiments in which NMR techniques have been used to implement the operations for a variety of quantum-information-processing jobs involving up to seven qubits (for reviews of NMR information processing, see Refs. [4–9]). The persistent skepticism has to do with questions about the “quantumness” of NMR information processing. Initially based on doubt that the highly mixed states used in NMR could be used to achieve genuinely quantum-mechanical effects, these questions were made concrete by the realization that all the quantum states accessed in present experiments are *unentangled* [10]. Entanglement is often thought to be an essential feature of quantum computation [11,12]. Arguments for an essential “quantumness” in NMR information processing are presented in Ref. [13], and an entirely different method for characterizing the “quantumness” of NMR is developed by Poulin [14].

The absence of entanglement in present NMR experiments means that the statistics of measurements made at any time during the experiments can be understood in terms of a

local realistic hidden-variable model [15] in which each spin has objective properties that determine the results of the measurements. In a local realistic model, the correlations observed in experiments can be attributed to classical correlations between realistic properties of the component qubits. The ability to describe the correlations observed in NMR experiments in terms of classical correlations between the qubits casts doubt on the “quantumness” of the experiments.

Modelling the statistics of the measurements made in NMR experiments is not sufficient for understanding the experiments. One must also be able to model the dynamics of the nuclear spins. NMR experimenters can implement with high accuracy any unitary operation, including the nonfactorizable unitary operations—those that cannot be written as a product of unitaries for each qubit—that produce entanglement when applied to pure quantum states. Previous attempts [16] to devise a local realistic description of the dynamics were only partially successful in that they did not provide a local realistic description of the changes produced by nonfactorizable unitaries which reproduced all the predictions of quantum mechanics. This left open the possibility that one might not be able to describe the correlations observed in successive measurements separated by nonfactorizable unitary operations in terms of local realistic properties and thus that present NMR experiments might violate *temporal* Bell inequalities [17] for successive measurements.

In this Letter we report a local realistic hidden-variable (LRHV) model for the states and dynamics of bulk-ensemble NMR information processing up to about 12 qubits. The existence of such a model rules out violation of any Bell-type inequality in present NMR experiments. This conclusion applies only to the bulk-ensemble model of information processing realized in present high-temperature, liquid-state NMR experiments; it does not apply to NMR methods based on distilling a pure state from a thermal state [5,18,19]. Our model is not satisfactory from the point of view developed by Schack and Caves [16] because the number of hidden variables scales exponentially with the number of qubits.

All NMR quantum computing experiments performed so far work in the following way. The state of each molecule, consisting of N active spin- $\frac{1}{2}$ nuclei, is described by a density operator

$$\hat{\rho} = (1 - \epsilon)\hat{1}/2^N + \epsilon\hat{\rho}_1, \quad (1)$$

which is a mixture of the desired state of the quantum computer, $\hat{\rho}_1$, with the maximally mixed state for N qubits, $\hat{1}/2^N$, $\hat{1}$ being the unit operator. When $\hat{\rho}_1$ is a pure state, $\hat{\rho}$ is called a *pseudopure state* [2].

The molecules in an NMR sample begin in thermal equilibrium, with a weak polarization $\alpha = h\nu/2kT \sim 2 \times 10^{-5}$ at room temperature, where $\nu \sim 300$ MHz is the average resonant frequency of the active spins in the strong longitudinal magnetic field. The first step in NMR information processing is to transform the molecules from equilibrium to a pseudopure state [3,20]. A consequence of pseudopure state synthesis is that the mixing parameter scales like $\epsilon = \alpha N/2^N$.

After synthesis of the desired initial state, the computation begins. The unitary operations required for the computation can be constructed from sequences of radio-frequency pulses alternating with periods of continuous evolution under the nuclear-spin Hamiltonian [4–9]. A unitary operator $\hat{U}$ takes an input state $\hat{\rho}$ to an output state

$$\hat{U}\hat{\rho}\hat{U}^\dagger = (1 - \epsilon)\hat{1}/2^N + \epsilon\hat{U}\hat{\rho}_1\hat{U}^\dagger. \quad (2)$$

The maximally mixed state is unaffected by the unitary transformation. The output state retains the form (1) with the same value of ϵ , and—this is the essence of the bulk-ensemble paradigm for quantum computation— $\hat{\rho}_1$ undergoes the desired unitary transformation.

The computation completed, the last step is to read out the answer. By applying radio-frequency pulses and then measuring the transverse magnetization of the sample, an NMR experimenter can determine the expectation value of any product of spin components, one for each qubit [4–9]. These expectation values have the form

$$\begin{aligned} C(\tilde{\mathbf{a}}) &\equiv C(\mathbf{a}_1, \dots, \mathbf{a}_N) \\ &= \text{tr}(\hat{\rho} \hat{\boldsymbol{\sigma}} \cdot \mathbf{a}_1 \otimes \dots \otimes \hat{\boldsymbol{\sigma}} \cdot \mathbf{a}_N) \\ &= \epsilon \text{tr}(\hat{\rho}_1 \hat{\boldsymbol{\sigma}} \cdot \mathbf{a}_1 \otimes \dots \otimes \hat{\boldsymbol{\sigma}} \cdot \mathbf{a}_N) . \end{aligned} \quad (3)$$

Here and throughout, a tilde over a quantity denotes a collection of N such quantities, one for each spin. In Eq. (3) the tensor product includes one operator for each spin; the vector operator $\hat{\boldsymbol{\sigma}} \equiv \hat{1}\mathbf{e}_0 + \hat{\sigma}_x\mathbf{e}_x + \hat{\sigma}_y\mathbf{e}_y + \hat{\sigma}_z\mathbf{e}_z$, where $\hat{\sigma}_x$, $\hat{\sigma}_y$, and $\hat{\sigma}_z$ are the Pauli operators; and $\mathbf{a}_r$ is either a spatial unit vector, in which case $\hat{\boldsymbol{\sigma}} \cdot \mathbf{a}_r$ is the component of spin r along the direction $\mathbf{a}_r$, or the unit vector $\mathbf{e}_0$ in the “zero” direction, in which case spin r does not contribute to the expectation value. The last equality in Eq. (3) assumes at least one of the vectors $\mathbf{a}_r$ is a spatial direction.

The expectation values (3) express the correlations between spin components of different spins. The maximally mixed state does not contribute to the correlation coefficients, which are determined by the state $\hat{\rho}_1$ that undergoes the desired evolution. The mixing parameter ϵ measures the strength of the magnetization signal. The scaling $\epsilon = \alpha N/2^N$ that comes from pseudopure state synthesis thus leads to an in-principle demand for an exponentially increasing number of molecules as the number of qubits increases [2,21], implying that bulk-ensemble quantum computation is not suitable for large-scale quantum computation.

Before constructing our LRHV model, we recall that any N -qubit density operator $\hat{\tau}$ has an associated quasidistribution [22]

$$w_{\hat{\tau}}(\tilde{\mathbf{n}}) \equiv \text{tr}(\hat{\tau} \hat{Q}(\tilde{\mathbf{n}})) , \quad (4)$$

where the vectors in the set $\tilde{\mathbf{n}} \equiv (\mathbf{n}_1, \dots, \mathbf{n}_N)$ are spatial unit vectors and

$$\hat{Q}(\tilde{\mathbf{n}}) \equiv \frac{1}{\mathcal{N}^N} (\hat{1} + 3\mathbf{n}_1 \cdot \hat{\boldsymbol{\sigma}}) \otimes \dots \otimes (\hat{1} + 3\mathbf{n}_N \cdot \hat{\boldsymbol{\sigma}}) . \quad (5)$$

For each spin, the unit vector $\mathbf{n}$ can point in $\mathcal{N}$ different directions satisfying

$$0 = \sum_{\mathbf{n}} n_j , \quad (6)$$

$$\frac{1}{3}\delta_{jk} = \frac{1}{\mathcal{N}} \sum_{\mathbf{n}} n_j n_k , \quad (7)$$

where the sums are over the possible directions and the subscripts indicate spatial components of $\mathbf{n}$. Condition (7) means that the vectors $\sqrt{3/\mathcal{N}}\mathbf{n}$ form a resolution of the 3-dimensional unit tensor; condition (6) places an additional constraint on the placement of the vectors. The vertices of a tetrahedron give the minimum number, $\mathcal{N} = 4$, of possible directions. The six vectors along the cardinal directions make up another simple possibility.

The density operator is then given by [22]

$$\hat{\tau} = \sum_{\tilde{\mathbf{n}}} w_{\hat{\tau}}(\tilde{\mathbf{n}}) |\tilde{\mathbf{n}}\rangle \langle \tilde{\mathbf{n}}| , \quad (8)$$

where $|\tilde{\mathbf{n}}\rangle \langle \tilde{\mathbf{n}}| \equiv |\mathbf{n}_1\rangle \langle \mathbf{n}_1| \otimes \cdots \otimes |\mathbf{n}_N\rangle \langle \mathbf{n}_N|$ and $|\mathbf{n}\rangle \langle \mathbf{n}| = \frac{1}{2}(\hat{1} + \hat{\boldsymbol{\sigma}} \cdot \mathbf{n})$ is the +1 eigenstate of $\hat{\boldsymbol{\sigma}} \cdot \mathbf{n}$. In terms of $w_{\hat{\tau}}(\tilde{\mathbf{n}})$, the correlation coefficients (3) take the form

$$C(\tilde{\mathbf{a}}) = \sum_{\tilde{\mathbf{n}}} w_{\hat{\tau}}(\tilde{\mathbf{n}}) (\mathbf{a}_1 \cdot \mathbf{m}_1) \cdots (\mathbf{a}_N \cdot \mathbf{m}_N) , \quad (9)$$

where $\mathbf{m}_r \equiv \mathbf{n}_r + \mathbf{e}_0$.

Under a unitary transformation the quasidistribution evolves according to

$$w_{\hat{U}\hat{\tau}\hat{U}^\dagger}(\tilde{\mathbf{n}}') = \sum_{\tilde{\mathbf{n}}} T_{\tilde{\mathbf{n}}'\tilde{\mathbf{n}}}^{\hat{U}} w_{\hat{\tau}}(\tilde{\mathbf{n}}) . \quad (10)$$

Here the transformation matrix $T^{\hat{U}}$ has matrix elements

$$T_{\tilde{\mathbf{n}}'\tilde{\mathbf{n}}}^{\hat{U}} \equiv \langle \tilde{\mathbf{n}} | \hat{U}^\dagger \hat{Q}(\tilde{\mathbf{n}}') \hat{U} | \tilde{\mathbf{n}} \rangle = w_{\hat{U}|\tilde{\mathbf{n}}\rangle \langle \tilde{\mathbf{n}}| \hat{U}^\dagger}(\tilde{\mathbf{n}}') , \quad (11)$$

given by the quasidistribution for $\hat{U}|\tilde{\mathbf{n}}\rangle$.

A *separable* density operator is one that has an ensemble decomposition in terms of product states. Such a state has no entanglement. If the quasidistribution $w_{\hat{\rho}}(\tilde{\mathbf{n}})$ is everywhere nonnegative, then $\hat{\rho}$ is definitely separable, and the statistics of all measurements can be understood in terms of classical tops whose probability to point in the directions $\tilde{\mathbf{n}}$ is $w_{\hat{\rho}}(\tilde{\mathbf{n}})$.

For any density operator, the quasidistribution satisfies $w_{\hat{\rho}}(\tilde{\mathbf{n}}) \geq [\text{minimum eigenvalue of } \hat{Q}(\tilde{\mathbf{n}})] = (-2)4^{N-1}/\mathcal{N}^N = -2^{2N-1}/\mathcal{N}^N$ [16,22]. Thus for density operators of the form (1), the quasidistribution is everywhere nonnegative if [10,22]

$$\epsilon \leq \frac{1}{1 + 2^{2N-1}} \equiv \eta . \quad (12)$$

Such states are *unentangleable* by any unitary transformation. For the polarization $\alpha \sim 2 \times 10^{-5}$ of present NMR experiments, all states up to about 12 qubits are unentangleable. It is known [23] that entangled states of the form (1) exist for $\epsilon > (1 + 2^{N-1})^{-1} \equiv \eta'$, i.e., $N \gtrsim 2/\alpha$, but whether there are entangled states for $\eta < \epsilon \leq \eta'$ is an open question.

We turn now to constructing a LRHV model for unentangleable states, i.e., for $\epsilon \leq \eta$. A straightforward model regards the directions $\tilde{\mathbf{n}}$ as hidden spin directions that determine the results of measurements stochastically. The problem with this simple model is that the change in the probability distribution $w_{\hat{\rho}}(\tilde{\mathbf{n}})$ as a consequence of a unitary transformation should be described by transition probabilities, which give the probability to go from initial directions $\tilde{\mathbf{n}}$ to final directions $\tilde{\mathbf{n}}'$. The matrix elements (11) provide candidate transition probabilities, but they cannot be interpreted as transition probabilities because they generally take on negative values. Schack and Caves [16] derived nonnegative transition probabilities from these matrix elements, but the dynamics described by these transition probabilities did not duplicate the predictions of quantum mechanics.

To include the unitary dynamics in the local realistic description, we construct a deterministic LRHV model that includes the probabilities $w_{\hat{\rho}}(\tilde{\mathbf{n}})$ in the set of hidden variables.

A quasidistribution $w_{\tilde{\mathbf{n}}}(\tilde{\mathbf{n}})$ can be regarded as a vector $\bar{w}_{\tilde{\mathbf{n}}}$ with $\mathcal{N}^N$ components labelled by the directions $\tilde{\mathbf{n}}$. An arbitrary vector in this space, denoted by $\bar{w}$, has components $w(\tilde{\mathbf{n}})$. We are only interested in normalized vectors, i.e., $\sum_{\tilde{\mathbf{n}}} w(\tilde{\mathbf{n}}) = 1$.

The hidden variables in our model are a vector $\bar{w}$, a set of spin directions $\tilde{\mathbf{n}}$, and a set of real variables $\tilde{\Lambda} \equiv (\Lambda_1, \dots, \Lambda_N)$ such that $-1 \leq \Lambda_r \leq 1$, $r = 1, \dots, N$. The hidden variables are denoted collectively by $\lambda = (\bar{w}, \tilde{\mathbf{n}}, \tilde{\Lambda})$. The probability density for λ is

$$P(\lambda) = \frac{1}{2^N} \delta(\bar{w} - \bar{w}_{\hat{\rho}}) w(\tilde{\mathbf{n}}) , \quad (13)$$

indicating that the variables $\tilde{\Lambda}$ are distributed randomly, the hidden vector $\bar{w}$ has a definite value given by $\bar{w}_{\hat{\rho}}$, and the probabilities for the hidden spin directions are determined by the hidden vector $\bar{w}$ and, hence, are written as $w(\tilde{\mathbf{n}})$. One can think of the hidden vector $\bar{w}$ as a set of parameters that weight a “roulette wheel,” so that when the wheel is “spun” to generate hidden spin directions, the probability for directions $\tilde{\mathbf{n}}$ is $w(\tilde{\mathbf{n}})$. Notice that the model requires that all components of $\bar{w}_{\hat{\rho}}$ be nonnegative, since these components become probabilities for the spin directions.

Measurement results are governed by functions $A_r(\mathbf{a}_r, \lambda)$, $r = 1, \dots, N$. The value of $A_r(\mathbf{a}_r, \lambda)$, either ± 1 , determines the result of a projective measurement of the component of spin r along spatial unit vector $\mathbf{a}_r$. The model is *realistic* because the results of spin-component measurements are determined by the hidden variables, and it is *local* because the result of a measurement of the component of spin r along direction $\mathbf{a}_r$ depends only on $\mathbf{a}_r$ and the hidden variables, not on measurement directions for other spins.

We choose the spin-component functions to be

$$A_r(\mathbf{a}_r, \lambda) = A_r(\mathbf{a}_r, \Lambda_r, \mathbf{n}_r) = \begin{cases} +1, & \text{if } \Lambda_r \geq -\mathbf{a}_r \cdot \mathbf{m}_r, \\ -1, & \text{if } \Lambda_r < -\mathbf{a}_r \cdot \mathbf{m}_r. \end{cases} \quad (14)$$

By using $\mathbf{m}_r \equiv \mathbf{n}_r + \mathbf{e}_0$, we can employ this function in cases where spin r is not involved in a measurement, i.e., when $\mathbf{a}_r = \mathbf{e}_0$, which gives $A_r(\mathbf{e}_0, \lambda) = 1$. The correlation coefficients predicted by the LRHV model,

$$\begin{aligned} C_{\text{LRHV}}(\tilde{\mathbf{a}}) &= \int d\lambda P(\lambda) A_1(\mathbf{a}_1, \lambda) \cdots A_N(\mathbf{a}_N, \lambda) \\ &= \sum_{\tilde{\mathbf{n}}} \int d\bar{w} \delta(\bar{w} - \bar{w}_{\hat{\rho}}) w(\tilde{\mathbf{n}}) \\ &\quad \times \prod_{r=1}^N \frac{1}{2} \int_{-1}^{+1} d\Lambda_r A_r(\mathbf{a}_r, \Lambda_r, \mathbf{n}_r) \\ &= \sum_{\tilde{\mathbf{n}}} w_{\hat{\rho}}(\tilde{\mathbf{n}}) (\mathbf{a}_1 \cdot \mathbf{m}_1) \cdots (\mathbf{a}_N \cdot \mathbf{m}_N) , \end{aligned} \quad (15)$$

duplicate the predictions of quantum mechanics.

Unitary dynamics fits easily into this model. For any unitary transformation $\hat{U}_{\tilde{\mathbf{z}}}$, the hidden variables for each molecule are updated in the following way: the variables $\tilde{\Lambda}$ are chosen randomly, the hidden vector $\bar{w}$ is updated deterministically using the transition matrix as in Eq. (10), and a new set of hidden spin directions is obtained by spinning a roulette wheel weighted by the new hidden vector, $\bar{w}_{\hat{\rho}\hat{U}^\dagger}$. By treating the components of

the hidden vector not as probabilities, but rather as parameters that are used to generate hidden spin directions stochastically, we avoid the need for a nonnegative transition matrix.

This description of the updating applies to a discrete unitary transformation, but it is easy to generalize it to a quasicontinuous hidden-variable dynamics: each molecule updates randomly in the fashion just described; i.e., each has a probability γdt to update within each time interval dt . The only requirement on this quasicontinuous dynamics is that the mean time γ^{-1} between updates be large compared to the precession time ν^{-1} of the nuclear spins in the strong magnetic field, but shorter than the duration of the radio-frequency pulses that are used to produce the desired dynamics.

It is trivial to generalize the LRHV model presented here to nonunitary evolutions, since these evolutions are, like unitary transformations, linear in the density operator. It is likely that the model could be extended to include all dynamics that accesses only separable states, i.e., states that have an expansion like Eq. (8), but with more general nonnegative quasidistributions than the canonical form (4).

The LRHV model developed here achieves our purpose of determining whether present NMR experiments can violate Bell inequalities. Bell inequalities are founded on two assumptions: the assumption that systems have objective properties and a “no-disturbance” assumption that asserts that the relevant measurements report faithfully the values of these properties. Standard Bell inequalities [15] justify the no-disturbance assumption from locality—measurements here cannot affect properties there. Temporal Bell inequalities [17], which involve successive measurements on a system, have a tougher time justifying the no-disturbance condition. Our LRHV model for NMR experiments describes the statistics of all measurements in terms of evolving classical correlations between realistic properties of the constituent nuclear spins. The conclusion is that NMR experiments up to about 12 qubits cannot violate any Bell inequality, temporal or otherwise.

Our purpose achieved, we acknowledge that our LRHV model is terribly contrived. It succeeds in giving a local realistic description of the dynamics by the brute force device of including an encoding of the entire density operator among the hidden variables. As a result, it requires an exponentially increasing number of hidden variables, $\sim 4^N$ in the most efficient version of the model. It leaves open the possibility that [16] the “quantumness” of NMR information processing lies in the ability to implement nonfactorizable unitaries that do not have an efficient local realistic description.

This work was partly supported by the National Security Agency (NSA) and the Advanced Research and Development Activity (ARDA) under Army Research Office (ARO) Contract No. DAAD19-01-1-0648. CMC received support from National Science Foundation Contract No. PHY99-07949 at the Institute for Theoretical Physics of the University of California, Santa Barbara.

REFERENCES

- [1] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, England, 2000).
- [2] D. G. Cory, A. F. Fahmy, and T. F. Havel, Proc. Natl. Acad. Sci. U.S.A. **94**, 1634 (1997).
- [3] N. A. Gershenfeld and I. L. Chuang, Science **275**, 350 (1997).
- [4] T. F. Havel, S. S. Somaroo, C.-H. Tseng, and D. G. Cory, Appl. Alg. Eng. Commun. Comp. **10**, 339 (2000).
- [5] D. G. Cory *et al.*, Fortschr. Phys. **48**, 875 (2000).
- [6] J. A. Jones, Fortschr. Phys. **48**, 909 (2000).
- [7] J. A. Jones, Prog. Nucl. Mag. Res. Spec. **38**, 325 (2001).
- [8] J. A. Jones, PhysChemComm **11**, 1 (2001).
- [9] L. M. K. Vandersypen, C. S. Yannoni, and I. L. Chuang, to be published in *Encyclopedia of Nuclear Magnetic Resonance*, edited by D. M. Grant and R. K. Harris (Wiley, New York, 2001), [arXiv.org e-print quant-ph/0012108](https://arxiv.org/abs/quant-ph/0012108).
- [10] S. L. Braunstein *et al.*, Phys. Rev. Lett. **83**, 1054 (1999).
- [11] R. Jozsa, *The Geometric Universe: Science, Geometry, and the Work of Roger Penrose*, edited by S. A. Huggett, L. J. Mason, K. P. Tod, S. T. Tsou, and N. M. J. Woodhouse (Oxford University, Oxford, England, 1998), p. 369.
- [12] A. Ekert and R. Jozsa, Phil. Trans. Roy. Soc. London A **356**, 1769 (1998). The authors argue that entanglement is the “essential non-classical feature providing the computational speed-up in the known quantum algorithms.”
- [13] R. Laflamme, D. G. Cory, C. Negrevergne, and L. Viola, unpublished, [arXiv.org e-print quant-ph/0110029](https://arxiv.org/abs/quant-ph/0110029).
- [14] D. Poulin, unpublished, [arXiv.org e-print quant-ph/0108102](https://arxiv.org/abs/quant-ph/0108102).
- [15] A. Peres, *Quantum Theory: Concepts and Methods* (Kluwer, Dordrecht, The Netherlands, 1993), Part II.
- [16] R. Schack and C. M. Caves, Phys. Rev. A **60**, 4354 (1999).
- [17] A. J. Leggett and A. Garg, Phys. Rev. Lett. **54**, 857 (1985); J. P. Paz and G. Mahler, Phys. Rev. Lett. **71**, 3235 (1993).
- [18] L. J. Schulman and U. Vazirani, in *Proceedings of the 31st Annual ACM Symposium on the Theory of Computation (STOC)* (ACM Press, 1999), p. 322; see also [arXiv.org e-print quant-ph/9804060](https://arxiv.org/abs/quant-ph/9804060).
- [19] E. Knill and R. Laflamme, Phys. Rev. Lett. **81**, 5672 (1998).
- [20] D. G. Cory, M. D. Price, and T. F. Havel, Physica D **120**, 82 (1998); I. L. Chuang, N. Gershenfeld, M. G. Kubinec, and D. W. Leung, Proc. Roy. Soc. London A **454**, 447 (1998); E. Knill, I. Chuang, and R. Laflamme, Phys. Rev. A **57**, 3348 (1998).
- [21] W. S. Warren, Science **277**, 1688 (1997).
- [22] R. Schack and C. M. Caves, J. Mod. Opt. **47**, 387 (2000).
- [23] W. Dür, J. I. Cirac, and R. Tarrach, Phys. Rev. Lett. **83**, 3562 (1999).